

Two-Photon and Two-gluon Decays of 0^{++} and 2^{++} P-wave Heavy Quarkonium States*

T. N. Pham

Centre de Physique Théorique, CNRS

Ecole Polytechnique,

91128 Palaiseau, Cedex, France

(Dated: October 4, 2010)

By neglecting the relative quark momenta in the propagator term, the two-photon and two-gluon decay amplitude of heavy quarkonia states can be written as a local heavy quark field operator matrix element which could be obtained from other processes or computed with QCD sum rules technique or lattice simulation, as shown in a recent work on $\eta_{c,b}$ two-photon decays. In this talk, I would like to discuss a similar calculation on P -wave $\chi_{c0,2}$ and $\chi_{b0,2}$ two-photon decays. We show that the effective Lagrangian for the two-photon decays of the P -wave $\chi_{c0,2}$ and $\chi_{b0,2}$ is given by the heavy quark energy-momentum tensor local operator and its trace, the $\bar{Q}Q$ scalar density. A simple expression for χ_{c0} two-photon and two-gluon decay rate in terms of the $f_{\chi_{c0}}$ decay constant, similar to that of η_c is obtained. From the existing QCD sum rules value for $f_{\chi_{c0}}$, we get 5 keV for the χ_{c0} two-photon width, somewhat larger than measurement.

PACS numbers: 13.20.Hd, 13.25.Gv, 11.10.St, 12.39.Hg

I. INTRODUCTION

First of all, I would like to dedicate this talk to the memory of Professor Giuseppe Nardulli, who, with great kindness and generosity has initiated the long and fruitful collaboration I have with the members of the Physics Department and INFN at the University of Bari.

In the non-relativistic bound state calculation [1, 2], the two-photon and two-gluon decay rates for P -wave quarkonium states depend on the derivative of the spatial wave function at the origin which has to be extracted from potential models, unlike the two-photon decay rate of S -wave η_c and η_b quarkonia which can be predicted from the corresponding J/ψ and Υ leptonic widths using heavy quark spin symmetry (HQSS) [3], there is no similar prediction for the P -wave χ_c and χ_b states and all the existing theoretical values for the decay rates are based on potential model

* Talk given at the *QCD@Work 2010 International Workshop on QCD: Theory and Experiment*, Martina Franca, Valle d'Istria, Italy, 20–23 June 2010

calculations [1, 4–14].

Since the matrix element of a heavy quark local operator between the vacuum and P -wave quarkonium state is also given, in bound state description, by the derivative of the spatial wave function at the origin, one could express the P -wave quarkonium two-photon and two-gluon decay amplitudes in terms of the matrix element of a local operator with the appropriate quantum number, like the heavy quark $\bar{Q}Q$ scalar density or axial vector current $\bar{Q}\gamma_\mu\gamma_5Q$. We have thus an effective Lagrangian for the two-photon and two-gluon decays of P -wave quarkonia in terms of heavy quark field operator instead of the traditional bound state description in terms of the wave function. This effective Lagrangian can be derived in a simple manner by neglecting the relative quark momentum in the heavy quark propagator as in non-relativistic bound state calculation. In this talk, I would like to report on a recent work [15] using the effective Lagrangian approach to describe the two-photon and two gluon decays of P -wave heavy quarkonia state, similar to that for S -wave quarkonia [3]. This was stimulated by the recent new CLEO measurements [16, 17] of the two-photon decay rates of the charmonium P -wave 0^{++} , χ_{c0} and 2^{++} χ_{c2} states. We obtain an effective Lagrangian for P -wave quarkonium decays in terms of the heavy quark energy-momentum tensor and its trace and that the two-photon and two-gluon decay rates of $\chi_{c0,2}$ and $\chi_{b0,2}$ can be expressed in terms of the decay constants $f_{\chi_{c0}}$ and $f_{\chi_{b0}}$, similar to that for η_c and η_b , which are given, respectively, by f_{η_c} and f_{η_b} . Then a calculation of $f_{\chi_{c0}}$ and $f_{\chi_{b0}}$ by sum rules technique [18, 19] or lattice simulation [20, 21] would give us a prediction of the P -wave quarkonia decay rates. In fact, as shown below, $f_{\chi_{c0}}$ obtained in [18] implies a value of 5 keV for the χ_{c0} two-photon width, somewhat larger than measurement. In the following I will present only the main results, as more details are given in the published paper [15].




FIG. 1: Diagrams for $Q\bar{Q}$ annihilation to two photons.

II. EFFECTIVE LAGRANGIAN FOR $\chi_{c0,2} \rightarrow \gamma\gamma$ AND $\chi_{b0,2} \rightarrow \gamma\gamma$

By neglecting term containing the relative quark momenta q in the quark propagator [25] ($Q_{c,b}^2$ being the heavy quark charge), the P -wave part of the $c\bar{c} \rightarrow \gamma\gamma, gg$ and $b\bar{b} \rightarrow \gamma\gamma, gg$ amplitudes

represented by diagrams in Fig. 1 are

$$\mathcal{M}(Q\bar{Q} \rightarrow \gamma\gamma) = -e^2 Q_{c,b}^2 \frac{A_{\mu\nu} \bar{v}(p_2) T_{\mu\nu} u(p_1)}{[(k_1 - k_2)^2/4 - m_Q^2]^2} \quad (1)$$

with $A_{\mu\nu}$ the photon part of the amplitude and the heavy quark part $T_{\mu\nu}$ given by

$$\begin{aligned} A_{\mu\nu} = & -2\epsilon_1 \cdot k_2 \epsilon_{2\mu} k_{1\nu} + 2\epsilon_1 \cdot \epsilon_2 k_{2\mu} k_{1\nu} \\ & -2\epsilon_2 \cdot k_1 \epsilon_{1\mu} k_{2\nu} + (k_1 \cdot k_2)(\epsilon_{1\mu} \epsilon_{2\nu} + \epsilon_{2\mu} \epsilon_{1\nu}) \end{aligned} \quad (2)$$

$$T_{\mu\nu} = (q_{1\mu} - q_{2\mu})\gamma_\nu \quad (3)$$

which can be obtained directly from the following effective Lagrangian for two-photon and two-gluon decay of P -wave heavy quarkonia states

$$\begin{aligned} \mathcal{L}_{\text{eff}}(Q\bar{Q} \rightarrow \gamma\gamma) &= -ic_1 A_{\mu\nu} \bar{Q}(\vec{\partial}_\mu - \overleftarrow{\partial}_\mu)\gamma_\nu Q \\ c_1 &= -e^2 Q_{c,b}^2 [(k_1 - k_2)^2/4 - m_Q^2]^{-2} \end{aligned} \quad (4)$$

With the matrix element of $\theta_{Q\mu\nu} = \bar{Q}(\vec{\partial}_\mu - \overleftarrow{\partial}_\mu)\gamma_\nu Q$ between the vacuum and $\chi_{c0,2}$ or $\chi_{b0,2}$ given by ($Q^2 = M^2$)

$$\begin{aligned} \langle 0 | \theta_{Q\mu\nu} | \chi_0 \rangle &= T_0 M^2 (-g_{\mu\nu} + Q_\mu Q_\nu / M^2), \\ \langle 0 | \theta_{Q\mu\nu} | \chi_2 \rangle &= -T_2 M^2 \epsilon_{\mu\nu}. \end{aligned} \quad (5)$$

The two-photon decay amplitudes are then easily obtained :

$$\mathcal{M}(\chi_0 \rightarrow \gamma\gamma) = -e^2 Q_{c,b}^2 \frac{T_0 A_0}{[M^2/4 + m_Q^2]^2} \quad (6)$$

$$\mathcal{M}(\chi_2 \rightarrow \gamma\gamma) = -e^2 Q_{c,b}^2 \frac{T_2 A_2}{[M^2/4 + m_Q^2]^2} \quad (7)$$

with $T_2 = \sqrt{3}T_0$ from HQSS and

$$A_0 = \left(\frac{3}{2}\right) M^2 (M^2 \epsilon_1 \cdot \epsilon_2 - 2\epsilon_1 \cdot k_2 \epsilon_2 \cdot k_1) \quad (8)$$

$$\begin{aligned} A_2 = & M^2 \epsilon_{\mu\nu} [M^2 \epsilon_{1\mu} \epsilon_{2\nu} - 2(\epsilon_1 \cdot k_2 \epsilon_{2\mu} k_{1\nu} + \epsilon_2 \cdot k_1 \epsilon_{1\mu} k_{2\nu} \\ & + \epsilon_1 \cdot \epsilon_2 k_{1\mu} k_{2\nu})] \end{aligned} \quad (9)$$

For QCD sum rules calculation or lattice simulation, it is simpler to compute the trace of the energy-momentum tensor $\theta_{Q\mu\mu}$ given by $2m_Q \bar{Q}Q$. We have then

$$\bar{v}(p_2) T_{\mu\mu} u(p_1) = 2m_Q \bar{v}(p_2) u(p_1) \quad (10)$$

The problem of computing the two-photon or two-gluon decay amplitude of $\chi_{c0,2}$ and $\chi_{b0,2}$ states is reduced to computing the decays constants $f_{\chi_{c0}}$ and $f_{\chi_{b0}}$ defined as

$$\langle 0 | \bar{Q}Q | \chi_0 \rangle = m_{\chi_0} f_{\chi_0} \quad (11)$$

Thus T_0 is given directly in terms of f_{χ_0} without using the derivative of the P -wave spatial wave function at the origin.

$$T_0 = \frac{f_{\chi_0}}{3} \quad (12)$$

Thus by comparing the expression for χ_{c0} and η_c we could already have some estimate for the χ_{c0} two-photon and two-gluon decay rates. For $f_{\chi_{c0}}$ of $O(f_{\eta_c})$, one would expect $\Gamma_{\gamma\gamma}(\chi_{c0})$ to be in the range of a few keV.

The decay rates of $\chi_{c0,2}$, $\chi_{b0,2}$ states can now be obtained in terms of the decay constant f_{χ_0} . We have :

$$\Gamma_{\gamma\gamma}(\chi_{c0}) = \frac{4\pi Q_c^4 \alpha_{em}^2 M_{\chi_{c0}}^3 f_{\chi_{c0}}^2}{(M_{\chi_{c0}} + b)^4} [1 + B_0(\alpha_s/\pi)], \quad (13)$$

$$\Gamma_{\gamma\gamma}(\chi_{c2}) = \left(\frac{4}{15}\right) \frac{4\pi Q_c^4 \alpha_{em}^2 M_{\chi_{c2}}^3 f_{\chi_{c0}}^2}{(M_{\chi_{c2}} + b)^4} [1 + B_2(\alpha_s/\pi)] \quad (14)$$

where $B_0 = \pi^2/3 - 28/9$ and $B_2 = -16/3$ are NLO QCD radiative corrections [22–24]

This expression is similar to that for η_c :

$$\Gamma_{\gamma\gamma}(\eta_c) = \frac{4\pi Q_c^4 \alpha_{em}^2 M_{\eta_c} f_{\eta_c}^2}{(M_{\eta_c} + b)^2} \left[1 - \frac{\alpha_s}{\pi} \frac{(20 - \pi^2)}{3} \right] \quad (15)$$

The two-gluon decay rates are :

$$\Gamma_{gg}(\chi_{c0}) = \left(\frac{2}{9}\right) \frac{4\pi \alpha_s^2 M_{\chi_{c0}}^3 f_{\chi_{c0}}^2}{(M_{\chi_{c0}} + b)^4} [1 + C_0(\alpha_s/\pi)], \quad (16)$$

$$\Gamma_{gg}(\chi_{c2}) = \left(\frac{4}{15}\right) \left(\frac{2}{9}\right) \frac{4\pi \alpha_s^2 M_{\chi_{c2}}^3 f_{\chi_{c0}}^2}{(M_{\chi_{c2}} + b)^4} [1 + C_2(\alpha_s/\pi)] \quad (17)$$

where $C_0 = 8.77$ and $C_2 = -4.827$ are NLO QCD radiative corrections. As with the two-photon decay rates, the expressions for two-gluon decay rates are similar to that for η_c :

$$\Gamma_{gg}(\eta_c) = \left(\frac{2}{9}\right) \frac{4\pi \alpha_s^2 M_{\eta_c} f_{\eta_c}^2}{(M_{\eta_c} + b)^2} \left[1 + 4.8 \frac{\alpha_s}{\pi} \right] \quad (18)$$

In a bound state calculation, using the relativistic spin projection operator [25, 26] , f_{η_c} and f_{χ_0} are given by

$$f_{\eta_c} = \sqrt{\frac{3}{32\pi m_Q^3}} \mathcal{R}_0(0) (4m_Q), \quad (19)$$

$$f_{\chi_0} = 12 \sqrt{\frac{3}{(8\pi m_Q)}} \left(\frac{\mathcal{R}'_1(0)}{M} \right) \quad (20)$$

$$(21)$$

which gives the decay amplitudes in agreement with the original calculation [1].

Comparing with f_{η_c} , we have

$$f_{\chi_{c0}} = 6 \left(\frac{\mathcal{R}'_1(0)}{\mathcal{R}_0(0)M} \right) f_{\eta_c}. \quad (22)$$

which becomes comparable to f_{η_c} .

Thus by comparing the expression for χ_{c0} and η_c we could already have some estimate for the χ_{c0} two-photon and two-gluon decay rates. For $f_{\chi_{c0}}$ of $O(f_{\eta_c})$, one would expect $\Gamma_{\gamma\gamma}(\chi_{c0})$ to be in the range of a few keV. As shown in Table 1, the predicted two-photon width of χ_{c0} from the sum rules value $f_{\chi_{c0}} = 357 \text{ MeV}$ [18] is however almost twice the CLEO value, but possibly with large theoretical uncertainties in sum rules calculation as to be expected, while a recent calculation [27] implies a larger decay rates for χ_{c0} . The measured ratio $\Gamma_{\gamma\gamma}(\chi_{c2})/\Gamma_{\gamma\gamma}(\chi_{c0})$ is then $\approx 0.24 \pm 0.09$, somewhat bigger than the predicted value of about 0.14 as shown in Table 1 together with the CLEO measurement of the decay rates [16] which gives $(2.53 \pm 0.37 \pm 0.26) \text{ keV}$ and $(0.60 \pm 0.06 \pm 0.06) \text{ keV}$ for χ_{c0} and χ_{c2} respectively.

Reference	$\Gamma_{\gamma\gamma}(\chi_{c0})(\text{keV})$	$\Gamma_{\gamma\gamma}(\chi_{c2})(\text{keV})$	$R = \frac{\Gamma_{\gamma\gamma}(\chi_{c2})}{\Gamma_{\gamma\gamma}(\chi_{c0})}$
Barbieri[1]	3.5	0.93	0.27
Godfrey[4]	1.29	0.46	0.36
Barnes[5]	1.56	0.56	0.36
Gupta[7]	6.38	0.57	0.09
Münz[8]	1.39 ± 0.16	0.44 ± 0.14	$0.32^{+0.16}_{-0.12}$
Huang[9]	3.72 ± 1.10	0.49 ± 0.16	$0.13^{+0.11}_{-0.06}$
Ebert[10]	2.90	0.50	0.17
Schuler[11]	2.50	0.28	0.11
Crater[12]	$3.34 - 3.96$	$0.43 - 0.74$	$0.13 - 0.19$
Wang[13]	3.78	—	—
Lavery[14]	$1.99 - 2.10$	$0.30 - 0.73$	$0.14 - 0.37$
This work	5.00	0.70	0.14
Exp(CLEO)[16]	$2.53 \pm 0.37 \pm 0.26$	$0.60 \pm 0.06 \pm 0.06$	$0.24 \pm 0.04 \pm 0.03$
Exp(Average)[16]	$2.31 \pm 0.10 \pm 0.12$	$0.51 \pm 0.02 \pm 0.02$	$0.20 \pm 0.01 \pm 0.02$

TABLE I: Potential model predictions for $\chi_{c0,2}$ two-photon widths compared with this work.

The two-photon $\chi_{c0,2}, \chi'_{c0,2}$ branching ratios are independent of $f_{\chi_{c0}}$

$$\mathcal{B}(\chi_{c0}, \chi'_{c0} \rightarrow \gamma\gamma) = \frac{9}{2} Q_c^4 \frac{\alpha_{em}^2}{\alpha_s^2} \left(1 + (B_0 - C_0) \frac{\alpha_s}{\pi} \right) \quad (23)$$

$$\mathcal{B}(\chi_{c2}, \chi'_{c2} \rightarrow \gamma\gamma) = \frac{6}{5} Q_c^4 \frac{\alpha_{em}^2}{\alpha_s^2} \left(1 + (B_2 - C_2) \frac{\alpha_s}{\pi} \right) \quad (24)$$

with $B_0 = \pi^2/3 - 28/9$, $B_2 = -16/3$, $C_0 = 8.77$, $C_2 = -4.827$. Apart from QCD radiative correction factors, the expressions for branching ratios are very similar to that for η_c and η'_c :

$$\mathcal{B}(\eta_c, \eta_{c'} \rightarrow \gamma\gamma) = \frac{9}{2} Q_c^4 \frac{\alpha_{em}^2}{\alpha_s^2} \left(1 - 8.2 \frac{\alpha_s}{\pi}\right) \quad (25)$$

with α_s evaluated at the appropriate scale.

For $\alpha_s = 0.26$, $\mathcal{B}(\eta_c \rightarrow \gamma\gamma) = 3.6 \times 10^{-4}$ to be compared with the measured value of $(2.8 \pm 0.9) \times 10^{-4}$ [17], but this prediction is rather sensitive to α_s , for example, with $\alpha_s = 0.28$, one would get $\mathcal{B}(\eta_c \rightarrow \gamma\gamma) = 2.95 \times 10^{-4}$, in better agreement with the measured value of $(2.4^{+1.1}_{-0.9}) \times 10^{-4}$ and for $\chi_{c0,2}$, the predicted two-photon branching ratios would be 3.45×10^{-4} and 4.45×10^{-4} compared with the measured values of $(2.35 \pm 0.23) \times 10^{-4}$ and $(2.43 \pm 0.18) \times 10^{-4}$, for χ_{c0} and χ_{c2} respectively. The predicted branching ratio for χ_{c2} is rather large and one would need $\alpha_s = 0.36$ to bring the predicted value closer to experiment.

Recently, the $Z(3930)$ state above $D\bar{D}$ threshold found by Belle [28] with mass $(3928 \pm 5 \pm 2)$ MeV and width $(29 \pm 10(\text{stat}) \pm 2(\text{sys}))$ MeV, consistent with χ'_{c2} , seems to be confirmed by the observation of a similar state by BaBar [29], with mass $(3926.7 \pm 2.7 \pm 1.1)$ MeV and width $(21.3 \pm 6.8 \pm 3.6)$ MeV. Belle [28] gives $\Gamma_{\gamma\gamma}(\chi'_{c2}) \times \mathcal{B}(D\bar{D}) = (0.18 \pm 0.05 \pm 0.03)$ keV while Babar [29] gives $\Gamma_{\gamma\gamma}(\chi'_{c2}) \times \mathcal{B}(D\bar{D}) = (0.24 \pm 0.05 \pm 0.04)$ keV for this state. If taken to be the $2P$ excited state χ'_{c2} and assuming $\mathcal{B}(D\bar{D}) \approx 0.70 - 1$ [30–32], one would get $\Gamma_{\gamma\gamma}(\chi'_{c2}) = (0.18 - 0.24 \pm 0.05 \pm 0.03)$ keV. This implies $f_{\chi'_{c0}} \simeq (195 - 225)$ MeV and $\Gamma_{gg}(\chi'_{c0})$ in the range $(5 - 10)$ MeV.

For $\chi_{b0,2}$ potential model calculations similar to that for $\chi_{c0,2}$, gives the two-photon width about 1/10 of that for η_b , which implies $f_{\chi_{b0}} = f_{\eta_b}/3$, smaller than Cornell potential [33] value $f_{\chi_{b0}} = 0.46 f_{\eta_b}$.

III. REMARK ON THE η'_c TWO-PHOTON DECAYS

Since the predicted two-photon branching ratios for $\chi_{c0,2}$, $\chi'_{c0,2}$ and for η_c, η'_c are similar and independent of the decay constants, apart from QCD radiative corrections, as seen in Eq. (23-24) and Eq. (25), one expects a large two-photon decay rates for η'_c , it would be relevant here to mention the problem of the $\eta'_c \rightarrow \gamma\gamma$ decay rate [3, 35]. The small value of $\Gamma_{\gamma\gamma}(\eta'_c) = (1.3 \pm 0.6)$ keV given previously by CLEO [34] is obtained by assuming $\mathcal{B}(\eta'_c \rightarrow K_S K \pi) \approx \mathcal{B}(\eta_c \rightarrow K_S K \pi)$. However, with the recent BaBar measurement of the ratio [36]

$$\begin{aligned} R(\eta_c(2S)K^+/\eta_c K^+) &= \frac{\mathcal{B}(B^+ \rightarrow \eta_c(2S)K^+) \times \mathcal{B}(\eta_c(2S) \rightarrow K\bar{K}\pi)}{\mathcal{B}(B^+ \rightarrow \eta_c K^+) \times \mathcal{B}(\eta_c \rightarrow K\bar{K}\pi)} \\ &= 0.096^{+0.020}_{-0.019}(\text{stat}) \pm 0.025(\text{syst}) \end{aligned} \quad (26)$$

and the Belle measurement [37]

$$\mathcal{B}(B^+ \rightarrow \eta_c K^+) \times \mathcal{B}(\eta_c \rightarrow K \bar{K} \pi) = (6.88 \pm 0.77_{-0.66}^{+0.55}) \times 10^{-5} \quad (27)$$

BABAR obtains [36]

$$\mathcal{B}(\eta'_c \rightarrow K_S K \pi) = (1.9 \pm 0.4(\text{stat}) \pm 1.1(\text{syst}))\%. \quad (28)$$

as quoted by CLEO [38]. This new BABAR value for $\mathcal{B}(\eta'_c \rightarrow K_S K \pi)$ is considerably smaller than the corresponding value $\mathcal{B}(\eta_c \rightarrow K_S K \pi) = (7.0 \pm 1.2)\%$ [17] for η_c .

Thus with the BaBar result for $\mathcal{B}(\eta'_c \rightarrow K_S K \pi)$ and the CLEO measurement [34]

$$R(\eta'_c/\eta_c) = \frac{\Gamma_{\gamma\gamma}(\eta'_c) \times \mathcal{B}(\eta'_c \rightarrow K_S K \pi)}{\Gamma_{\gamma\gamma}(\eta_c) \times \mathcal{B}(\eta_c \rightarrow K_S K \pi)} = 0.18 \pm 0.05 \pm 0.02 \quad (29)$$

one would get [38]

$$\Gamma(\eta_{c'} \rightarrow \gamma\gamma) = (4.8 \pm 3.7) \text{ keV} \quad (30)$$

in agreement with the predicted value

$$\Gamma(\eta_{c'} \rightarrow \gamma\gamma) = (4.1 \pm 2.3) \text{ keV} \quad (31)$$

while the assumption of near equality of the $K_S K \pi$ branching ratios for η_c and η'_c

$$\mathcal{B}(\eta'_c \rightarrow K_S K \pi) \approx \mathcal{B}(\eta_c \rightarrow K_S K \pi) \quad (32)$$

and the Belle ratio [39]

$$R(\eta'_c K/\eta_c K) = \frac{\mathcal{B}(B \rightarrow K \eta_c(2S) \times \mathcal{B}(\eta_c(2S) \rightarrow K_S K^- \pi^+)}{\mathcal{B}(B^0 \rightarrow K \eta_c) \times \mathcal{B}(\eta_c \rightarrow K_S K^- \pi^+)} = 0.38 \pm 0.12 \pm 0.05 \quad (33)$$

would lead to [38]

$$\Gamma_{\gamma\gamma}(\eta'_c) = (1.3 \pm 0.6) \text{ keV} \quad (34)$$

which is rather small compared with the predicted value given in Eq. (31) above.

IV. CONCLUSION

In conclusion, we have derived an effective Lagrangian for $\chi_{c0,2}$ and $\chi_{b0,2}$ two-photon and two-gluon in terms of the decay constants $f_{\chi_{c,b0}}$, similar to that for $\eta_{c,b}$ in terms of $f_{\eta_{c,b}}$.

Existing sum rules calculation, however produces a two-photon width about 5 keV, somewhat bigger than the CLEO measured value. It remains to be seen whether a better determination of $f_{\chi_{c0}}$ from lattice simulation or QCD sum rules calculation could bring the $\chi_{c0,2}$ two-photon decay rates closer to experiments or higher order QCD radiative corrections and large relativistic corrections are needed to explain the data.

The problem of two-photon width of η'_c would go away if more data could confirm the small BaBar value for $\mathcal{B}(\eta'_c \rightarrow K_S K \pi)$ compared with $\mathcal{B}(\eta_c \rightarrow K_S K \pi)$.

As relativistic corrections should be small for P -wave bottomia $\chi_{b0,2}$ states, two-photon and two-gluon decays could provide a test of QCD and a determination of α_s at the m_b mass scale.

V. ACKNOWLEDGMENTS

I would like to thank P. Colangelo, F. De Fazio and the organizers for generous support and warm hospitality extended to me at Martina Franca. This work was supported in part by the EU contract No. MRTN-CT-2006-035482, "FLAVIANet".

-
- [1] R. Barbieri, R. Gatto and R. Kogerler, Phys. Lett. B **60**, 183 (1976).
 - [2] For a review, see N. Brambilla *et al.*, *Heavy quarkonium physics*, CERN Yellow Report, CERN-2005-005, 2005.
 - [3] J. P. Lansberg and T. N. Pham, Phys. Rev. D **74**, 034001 (2006); *ibid* D **75**, 017501 (2007).
 - [4] S. Godfrey and N. Isgur, Phys. Rev. D **32**, 189 (1985).
 - [5] T. Barnes, *Proceedings of the IX International Workshop on Photon-Photon Collisions*, edited by D. O. Caldwell and H. P. Paar (World Scientific, Singapore, 1992) p. 263.
 - [6] G. Bodwin, E. Braaten, and G. P. Lepage, Phys. Rev. D **46**, R1914 (1992).
 - [7] S. N. Gupta, J. M. Johnson, and W. W. Repko, Phys. Rev. D **54**, 2075 (1996);
 - [8] C. R. Münz, Nucl. Phys. A **609**, 364 (1996).
 - [9] H.-W. Huang and K.-Ta Chao, Phys. Rev. D **54**, 6850 (1996); *errata* Phys. Rev. D **56** 1821 (1996).
 - [10] D. Ebert, R. N. Faustov and V. O. Galkin, Mod. Phys. Lett. A **18**, (2003).
 - [11] G. A. Schuler, F. A. Berends, and R. van Gulik, Nucl. Phys. B **523**, 423 (1998).
 - [12] H. W. Crater, C. Y. Wong and P. Van Alstine, Phys. Rev. D **74**, 054028 (2006).

- [13] G.-Li Wang, Phys. Lett. B **653**, 206 (2007).
- [14] J. T. Lavery, S. F. Radford, and W. W. Repko, [arXiv:0901.3917] [hep-ph].
- [15] J. P. Lansberg and T. N. Pham, Phys. Rev. D **79**, 094016 (2009).
- [16] K. M. Ecklund *et al* (CLEO Collaboration), Phys. Rev. D **78**, 091501 (2008) and other recent results quoted therein.
- [17] C. Amsler, *et al*, Particle Data Group, *Review of Particle Physics*, Phys. Lett. B **667**, 1 (2008).
- [18] V. A. Novikov, L. B. Okun, M. A. Shifman, A. I. Vainshtein, M. B. Voloshin and V. I. Zakharov, Phys. Rept. **41**, 1 (1978).
- [19] L. J. Reinders, H. R. Rubinstein and S. Yazaki, Phys. Lett. B **113**, 411 (1982).
- [20] J. J. Dudek, R. G. Edwards and D. G. Richards, Phys. Rev. D **73**, 074507 (2006).
- [21] T. W. Chiu, T. H. Hsieh, and Ogawa [TWQCD Collaboration], Phys. Lett. B **651**, 171 (2007).
- [22] R. Barbieri, M. Caffo, R. Gatto and E. Remiddi, Phys. Lett. B **95**, 93 (1980);
Nucl. Phys. B **192**, 61 (1981).
- [23] W. Kwong, P. B. Mackenzie, R. Rosenfeld and J. L. Rosner, Phys. Rev. D **37**, 3210 (1988).
- [24] M. Mangano and A. Petrelli, Phys. Lett. B **352**, 445 (1995).
- [25] J. H. Kühn, J. Kaplan and E. G. O. Safiani, Nucl. Phys. B **157**, 125 (1979).
- [26] B. Guberina, J. H. Kühn, R. D. Peccei, and R. Ruckl, Nucl. Phys. B **174**, 317 (1980).
- [27] P. Colangelo, F. De Fazio, and T.N. Pham, Phys. Lett. B **542**, 71 (2002).
- [28] S. Uehara *et al.* (BELLE collaboration), Phys. Rev. D **96**, 082003 (2006).
- [29] B. Aubert *et al.* (BABAR Collaboration), Phys. Rev. D **81**, 092003 (2010).
- [30] P. Colangelo, Private Communication.
- [31] O. Lakhina and E. S. Swanson, Phys. Rev. D **74**, 014012 (2006).
- [32] B-Q. Li and K. T. Chao, [arXiv:0903.5506] [hep-ph].
- [33] E. Eichten and C. Quigg, Phys. Rev. D **52**, 1726 (1995).
- [34] D. M. Asner *et al.* (CLEO Collaboration)], Phys. Rev. Lett. **92**, 142001 (2004).
- [35] T. N. Pham, Proceedings of the International Workshop on Quantum Chromodynamics Theory and Experiment, Martina Franca, Valle d'Itria, Italy, 16-20 June 2007, AIP Conf. Proc. **964**, 124, (2007);
J. P. Lansberg and T. N. Pham, Proceedings of the Joint Meeting Heidelberg-Liège-Paris-Wroclaw Hadronic Physics HLPW 2008, Spa Belgium, 6-8 March 2008, AIP Conf. Proc. **1038**, 259 (2008).
- [36] B. Aubert *et al.* (BABAR Collaboration), Phys. Rev. D **78**, 012006 (2008).
- [37] F. Fang *et al.* (BELLE collaboration), Phys. Rev. Lett. **90**, 071801 (2003).
- [38] D. Cronin-Hennessy *et al.* (CLEO Collaboration), Phys. Rev. D **81**, 052002 (2010).
- [39] S. K. Choi *et al.* (BELLE collaboration), Phys. Rev. Lett. **89**, 102001 (2002).